

9. The z -transform

- the z -transform
- finding inverse z -transform
- properties of z -transform

Definition

the z -transform of the sequence $x[n]$ is

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$


- where z is a complex variable
- the *region of convergence* (ROC) for $X(z)$ are the values of z (the region in the complex plane) for which the sum converges (or exists)

Example: the z -transform for the signal $\gamma^n u[n]$ is

$$X(z) = \sum_{n=0}^{\infty} \gamma^n u[n] z^{-n} = \sum_{n=0}^{\infty} \left(\frac{\gamma}{z}\right)^n = \frac{1}{1 - \frac{\gamma}{z}} = \frac{z}{z - \gamma} \quad \left| \frac{\gamma}{z} \right| < 1 \quad (|z| > |\gamma|)$$

for $|z| < |\gamma|$, the sum does not converge (it goes to infinity)

the ROC of $X(z)$ is the shaded region outside the circle of radius $|\gamma|$ (in z -plane)



Uniqueness for causal signals

- the z -transform of the z -transform of $-\gamma^n u[-(n+1)]$ is also $z/(z-\gamma)$ (but different ROC $|z| < |\gamma|$)
- the inverse z -transform of $z/(z-\gamma)$ is not unique
- restricting the $x[n]$ to be causal, then the inverse transform is unique

The unilateral z -transform

the *unilateral z -transform* is defined for causal signals

$$X(z) = \sum_{n=0}^{\infty} x[n]z^{-n}$$

Existence: the existence of the z -transform is guaranteed if

$$|X(z)| \leq \sum_{n=0}^{\infty} \frac{|x[n]|}{|z|^n} < \infty, \quad \text{for some } |z|$$

- if $|x[n]| \leq r_0^n$ for some r_0 , then

$$|X(z)| \leq \sum_{n=0}^{\infty} \left(\frac{r_0}{|z|} \right)^n = \frac{1}{1 - \frac{r_0}{|z|}}, \quad |z| > r_0$$


and hence, $X(z)$ exists for $|z| > r_0$

- some signal models (e.g., γ^{n^2} grow faster than the exponential signal r_0^n (for any r_0) and therefore are not z -transformable

Example 9.1

using the definition of the z -transform, find the z -transforms of:

- (a) $\delta[n]$
- (b) $u[n]$
- (c) $\cos(\beta n)u[n]$
- (d) the signal $x[n]$ shown below



Solution:

(a)

$$X(z) = \sum_{n=0}^{\infty} x[n]z^{-n} = \sum_{n=0}^{\infty} \delta[n]z^{-n} = 1$$

therefore,

$$\delta[n] \iff 1 \quad \text{for all } z$$

(b) for $x[n] = u[n]$, we have

$$\begin{aligned} X(z) &= \sum_{n=0}^{\infty} x[n]z^{-n} = \sum_{n=0}^{\infty} z^{-n} \\ &= \frac{1}{1 - \frac{1}{z}} \quad \left| \frac{1}{z} \right| < 1 \\ &= \frac{z}{z-1} \quad |z| > 1 \end{aligned}$$

therefore,

$$u[n] \iff \frac{z}{z-1} \quad |z| > 1$$

(c) using $\cos(\beta n) = (e^{j\beta n} + e^{-j\beta n}) / 2$ and

$$e^{\pm j\beta n} u[n] \iff \frac{z}{z - e^{\pm j\beta}} \quad |z| > |e^{\pm j\beta}| = 1$$

we have

$$X(z) = \frac{1}{2} \left[\frac{z}{z - e^{j\beta}} + \frac{z}{z - e^{-j\beta}} \right] = \frac{z(z - \cos \beta)}{z^2 - 2z \cos \beta + 1} \quad |z| > 1$$

(d) here, we have

$$X(z) = 1 + \frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} + \frac{1}{z^4} = \frac{z^4 + z^3 + z^2 + z + 1}{z^4} \quad \text{for all } z \neq 0$$

or

$$X(z) = \sum_{n=0}^4 z^{-n} = \frac{\left(\frac{1}{z}\right)^5 - \left(\frac{1}{z}\right)^0}{\frac{1}{z} - 1} = \frac{z}{z-1} (1 - z^{-5})$$

Exercises

- find the z -transform of



Answer: $X(z) = \frac{z^5 + z^4 + z^3 + z^2 + z + 1}{z^9}$ or $\frac{z}{z-1} (z^{-4} - z^{-10})$

- use the table to find the z -transform of

$$x[n] = 20.65(\sqrt{2})^n \cos[(\pi/4)n - 1.415]u[n]$$

Answer: $\frac{z(3.2z + 17.2)}{z^2 - 2z + 2}$

Outline

- the z -transform
- **finding inverse z -transform**
- properties of z -transform

Inverse z -transform

the *inverse z -transform* of $X(z)$ is:

$$x[n] = \frac{1}{2\pi j} \oint X(z)z^{n-1}dz$$

- the symbol \oint indicates an integration in counterclockwise direction around a closed path in the complex plane
- we shall avoid the integration in the complex plane required to find the inverse z -transform by using the (unilateral) transform table

Inverse transform using known z -transform pairs

- many of the transforms $X(z)$ of practical interest are rational functions $X(z) = A(z)/B(z)$ (ratio of polynomials in z)
- if we can express $X(z)$ as a sum of partial fractions, then we can find the inverse transforms using the z -transform table

Notation

$$X(z) = \mathcal{Z}\{x[n]\} \quad \text{and} \quad x[n] = \mathcal{Z}^{-1}\{X(z)\}$$

$$x[n] \iff X(z)$$

note that

$$\mathcal{Z}^{-1}[\mathcal{Z}\{x[n]\}] = x[n] \quad \text{and} \quad \mathcal{Z}[\mathcal{Z}^{-1}\{X(z)\}] = X(z)$$

Linearity: if

$$x_1[n] \iff X_1(z) \quad \text{and} \quad x_2[n] \iff X_2(z)$$

then

$$a_1x_1[n] + a_2x_2[n] \iff a_1X_1(z) + a_2X_2(z)$$

Example 9.2

find the inverse z -transforms of

$$\frac{8z - 19}{(z - 2)(z - 3)}$$

Solution: expanding $X(z)$ into partial fractions yields

$$X(z) = \frac{8z - 19}{(z - 2)(z - 3)} = \frac{3}{z - 2} + \frac{5}{z - 3}$$

from table (pair 7), we obtain

$$x[n] = [3(2)^{n-1} + 5(3)^{n-1}] u[n - 1]$$

Modified partial fraction expansion: to obtain a form that contains $u[n]$ rather than $u[n - 1]$, we can instead expand:

$$\frac{X(z)}{z} = \frac{8z - 19}{z(z - 2)(z - 3)} = \frac{(-19/6)}{z} + \frac{(3/2)}{z - 2} + \frac{(5/3)}{z - 3}$$

multiplying both sides by z yields

$$X(z) = -\frac{19}{6} + \frac{3}{2} \left(\frac{z}{z - 2} \right) + \frac{5}{3} \left(\frac{z}{z - 3} \right)$$

using pairs 1 and 6 in table, we get

$$x[n] = -\frac{19}{6} \delta[n] + \left[\frac{3}{2} (2)^n + \frac{5}{3} (3)^n \right] u[n]$$

we often expand $X(z)/z$ rather than $X(z)$ into partial fractions and then multiply both sides by z to obtain modified partial fractions of $X(z)$

Example 9.3

find the inverse z -transforms of

$$(a) \frac{z(2z^2 - 11z + 12)}{(z - 1)(z - 2)^3}$$

$$(b) \frac{2z(3z + 17)}{(z - 1)(z^2 - 6z + 25)}$$

Solution:

(a) we have

$$\frac{X(z)}{z} = \frac{2z^2 - 11z + 12}{(z-1)(z-2)^3} = \frac{k}{z-1} + \frac{a_0}{(z-2)^3} + \frac{a_1}{(z-2)^2} + \frac{a_2}{(z-2)}$$

where

$$k = \left. \frac{2z^2 - 11z + 12}{(z-1)(z-2)^3} \right|_{z=1} = -3 \quad a_0 = \left. \frac{2z^2 - 11z + 12}{(z-1)(z-2)^3} \right|_{z=2} = -2$$

therefore,

$$\frac{X(z)}{z} = \frac{2z^2 - 11z + 12}{(z-1)(z-2)^3} = \frac{-3}{z-1} - \frac{2}{(z-2)^3} + \frac{a_1}{(z-2)^2} + \frac{a_2}{(z-2)}$$

to find a_2 , we multiply both sides by z and let $z \rightarrow \infty$:

$$0 = -3 - 0 + 0 + a_2 \implies a_2 = 3$$

letting z take any convenient value, say, $z = 0$, on both sides:

$$\frac{12}{8} = 3 + \frac{1}{4} + \frac{a_1}{4} - \frac{3}{2}$$

which yields $a_1 = -1$; therefore,

$$\frac{X(z)}{z} = \frac{-3}{z-1} - \frac{2}{(z-2)^3} - \frac{1}{(z-2)^2} + \frac{3}{z-2}$$

and

$$X(z) = -3\frac{z}{z-1} - 2\frac{z}{(z-2)^3} - \frac{z}{(z-2)^2} + 3\frac{z}{z-2}$$

using pairs 6 and 10 of table, gives

$$\begin{aligned} x[n] &= \left[-3 - 2\frac{n(n-1)}{8}(2)^n - \frac{n}{2}(2)^n + 3(2)^n \right] u[n] \\ &= -\left[3 + \frac{1}{4}(n^2 + n - 12)2^n \right] u[n] \end{aligned}$$

(b) **complex poles**

$$X(z) = \frac{2z(3z + 17)}{(z - 1)(z^2 - 6z + 25)} = \frac{2z(3z + 17)}{(z - 1)(z - 3 - j4)(z - 3 + j4)}$$

method of first-order factors:

$$\frac{X(z)}{z} = \frac{2(3z + 17)}{(z - 1)(z^2 - 6z + 25)} = \frac{2(3z + 17)}{(z - 1)(z - 3 - j4)(z - 3 + j4)}$$

we find the partial fraction of $X(z)/z$ using the Heaviside "cover-up" method:

$$\frac{X(z)}{z} = \frac{2}{z - 1} + \frac{1.6e^{-j2.246}}{z - 3 - j4} + \frac{1.6e^{j2.246}}{z - 3 + j4}$$

and

$$X(z) = 2\frac{z}{z - 1} + (1.6e^{-j2.246})\frac{z}{z - 3 - j4} + (1.6e^{j2.246})\frac{z}{z - 3 + j4}$$

the inverse transform of the complex two terms (complex conjugate poles) can be obtained from pair 12b (table) with $r/2 = 1.6$, $\theta = -2.246$ rad, $\gamma = 3 + j4 = 5e^{j0.927}$, so that $|\gamma| = 5$, $\beta = 0.927$; therefore, we have

$$x[n] = [2 + 3.2(5)^n \cos(0.927n - 2.246)] u[n]$$

method of quadratic factors:

$$\frac{X(z)}{z} = \frac{2(3z + 17)}{(z - 1)(z^2 - 6z + 25)} = \frac{2}{z - 1} + \frac{Az + B}{z^2 - 6z + 25}$$

multiplying both sides by z and letting $z \rightarrow \infty$, we find

$$0 = 2 + A \implies A = -2$$

and

$$\frac{2(3z + 17)}{(z - 1)(z^2 - 6z + 25)} = \frac{2}{z - 1} + \frac{-2z + B}{z^2 - 6z + 25}$$

to find B , we let z take any convenient value, say, $z = 0$:

$$\frac{-34}{25} = -2 + \frac{B}{25} \implies B = 16$$

therefore,

$$\frac{X(z)}{z} = \frac{2}{z-1} + \frac{-2z+16}{z^2-6z+25}$$

and

$$X(z) = \frac{2z}{z-1} + \frac{z(-2z+16)}{z^2-6z+25}$$

we now use pair 12c (table) with $A = -2$, $B = 16$, $|\gamma| = 5$, and $a = -3$, so that

$$r = \sqrt{\frac{100 + 256 - 192}{25 - 9}} = 3.2, \quad \beta = \cos^{-1}\left(\frac{3}{5}\right) = 0.927 \text{ rad}$$

and

$$\theta = \tan^{-1}\left(\frac{-10}{-8}\right) = -2.246 \text{ rad}$$

hence

$$x[n] = [2 + 3.2(5)^n \cos(0.927n - 2.246)] u[n]$$

Inverse transform by power series expansion

by definition,

$$X(z) = \sum_{n=0}^{\infty} x[n]z^{-n} = x[0]z^0 + x[1]z^{-1} + x[2]z^{-2} + x[3]z^{-3} + \dots$$

- if we can expand $X(z)$ into the power series in z^{-1} , the coefficients of this power series can be identified as $x[0], x[1], x[2], x[3], \dots$
- a rational $X(z)$ can be expanded into a power series of z^{-1} by dividing its numerator by the denominator
- this procedure it is only useful if we want to know only the first few terms of the sequence $x[n]$

Example:

$$X(z) = \frac{z^2(7z - 2)}{(z - 0.2)(z - 0.5)(z - 1)} = \frac{7z^3 - 2z^2}{z^3 - 1.7z^2 + 0.8z - 0.1}$$

we have:

$$X(z) = \frac{z^2(7z - 2)}{(z - 0.2)(z - 0.5)(z - 1)} = 7 + 9.9z^{-1} + 11.23z^{-2} + 11.87z^{-3} + \dots$$

therefore,

$$x[0] = 7, \quad x[1] = 9.9, \quad x[2] = 11.23, \quad x[3] = 11.87, \quad \dots$$

Exercises

- using long division to find the power series in z^{-1} , show that the inverse z -transform of $z/(z - 0.5)$ is $(0.5)^n u[n]$ or $2^{-n} u[n]$
- find the inverse z -transform of the following functions:

$$(a) \frac{z(2z - 1)}{(z - 1)(z + 0.5)}$$

$$(b) \frac{1}{(z - 1)(z + 0.5)}$$

$$(c) \frac{1}{(z + 2)(z - 0.5)^2}$$

$$(d) \frac{5z(z - 1)}{z^2 - 1.6z + 0.8} \quad [\text{Hint: } \sqrt{0.8} = 2/\sqrt{5}]$$

Answers:

$$(a) \left[\frac{2}{3} + \frac{4}{3}(-0.5)^n \right] u[n]$$

$$(b) -2\delta[n] + \left[\frac{2}{3} + \frac{4}{3}(-0.5)^n \right] u[n]$$

$$(c) 18\delta[n] - [0.72(-2)^n + 17.28(0.5)^n - 14.4n(0.5)^n] u[n]$$


$$(d) \frac{5\sqrt{5}}{2} \left(\frac{2}{\sqrt{5}} \right)^n \cos(0.464n + 0.464) u[n]$$

Outline


- the z -transform
- finding inverse z -transform
- **properties of z -transform**

Shifting forms


we need to distinguish between: $x[n]u[n]$, $x[n - m]u[n]$, $x[n - m]u[n - m]$,
and $x[n + m]u[n]$




(a)




(b)



(d)



(c)



(e)

Shifting properties

Causal-part right-shift: if $x[n]u[n] \iff X(z)$ then for integer value of m ,

$$x[n-m]u[n-m] \iff \frac{1}{z^m}X(z)$$

Right-shift: integer value of $m > 0$,

$$x[n-m]u[n] \iff z^{-m}X(z) + z^{-m} \sum_{n=1}^m x[-n]z^n$$

for $m = 1$:

$$x[n-1]u[n] \iff \frac{1}{z}X(z) + x[-1]$$

Left-shift: for integer value of $m > 0$,

$$x[n+m]u[n] \iff z^mX(z) - z^m \sum_{n=0}^{m-1} x[n]z^{-n}$$

for $m = 1$:

$$x[n+1]u[n] \iff zX(z) - zx[0]$$

Left-shift (advance)

if $x[n]u[n] \iff X(z)$, then


$$x[n+1]u[n] \iff zX(z) - zx[0]$$

for integer value of $m > 0$,

$$x[n+m]u[n] \iff z^m X(z) - z^m \sum_{n=0}^{m-1} x[n]z^{-n}$$

Example 9.4

use the shifting property and the z -transform table to find the z -transform of $x[n]$



Solution: $x[n]$ can be expressed as:

$$x[n] = n(u[n] - u[n - 6]) = nu[n] - nu[n - 6]$$

to use the right-shift property, we rearrange $nu[n-6]$ in terms of $(n-6)u[n-6]$:

$$\begin{aligned}x[n] &= nu[n] - (n-6+6)u[n-6] \\ &= nu[n] - (n-6)u[n-6] - 6u[n-6]\end{aligned}$$

because $u[n] \iff z/(z-1)$,

$$u[n-6] \iff \frac{1}{z^6} \frac{z}{z-1} = \frac{1}{z^5(z-1)}$$

also, because $nu[n] \iff z/(z-1)^2$

$$(n-6)u[n-6] \iff \frac{1}{z^6} \frac{z}{(z-1)^2} = \frac{1}{z^5(z-1)^2}$$

therefore,

$$X(z) = \frac{z}{(z-1)^2} - \frac{1}{z^5(z-1)^2} - \frac{6}{z^5(z-1)} = \frac{z^6 - 6z + 5}{z^5(z-1)^2}$$

Time-reversal

Time-reversal: if $x[n] \iff X(z)$, then

$$x[-n] \iff X(1/z)$$

- the region of convergence is also inverted
- for example, if the ROC of $x[n]$ is $|z| > |\gamma|$, then the ROC of $x[-n]$ is $|z| < 1/|\gamma|$

Example: find the z -transform and ROC of $u[-n]$

from table $u[n] \iff U(z) = z/(z-1)$ with ROC $|z| > 1$, hence from time-reversal property, we have

$$u[-n] \iff U(1/z) = \frac{1/z}{(1/z) - 1} = \frac{1}{z-1}$$

with inverted ROC $|z| < 1$

Scaling and differentiation in the z -domain

z -domain scaling (multiplication by γ^n): if $x[n]u[n] \iff X(z)$, then

$$\gamma^n x[n]u[n] \iff X\left(\frac{z}{\gamma}\right)$$

z -domain differentiation (multiplication by n): if $x[n]u[n] \iff X(z)$, then

$$nx[n]u[n] \iff -z \frac{d}{dz} X(z)$$

Example 9.5

find the unilateral z -transform of $x[n] = (1 - n) \cos(\frac{\pi}{2}(n - 1))u[n - 1]$

Solution: using properties:

$$\cos(\pi n/2)u[n] \iff \frac{z^2}{z^2 + 1} \quad (\text{table pair 11a})$$

$$-n \cos(\pi n/2)u[n] \iff z \frac{d}{dz} \left(\frac{z^2}{z^2 + 1} \right) \quad (z\text{-domain differentiation})$$

$$= z \left(\frac{2z}{z^2 + 1} - \frac{z^2}{(z^2 + 1)^2} (2z) \right) = \frac{2z^2}{(z^2 + 1)^2}$$

$$x[n] = -(n - 1) \cos(\pi(n - 1)/2)u[n - 1] \iff z^{-1} \frac{2z^2}{(z^2 + 1)^2} \quad (\text{time shift})$$

therefore,

$$X(z) = \frac{2z}{(z^2 + 1)^2} = \frac{2z}{z^4 + 2z^2 + 1}$$

Convolution

Time-convolution: if

$$x_1[n] \iff X_1(z) \quad \text{and} \quad x_2[n] \iff X_2(z)$$

then

$$x_1[n] * x_2[n] \iff X_1(z)X_2(z)$$

Example: using z -transform compute $u[n] * u[n - 1]$

from convolution and shifting properties, we have

$$u[n] * u[n - 1] \iff \frac{z}{z - 1} \frac{z}{z - 1} \frac{1}{z} = \frac{z}{(z - 1)^2}$$

using the table, we have

$$u[n] * u[n - 1] = nu[n] \iff \frac{z}{(z - 1)^2}$$

Initial and final values

Initial value theorem: for a causal $x[n]$, we have


$$x[0] = \lim_{z \rightarrow \infty} X(z)$$

Final value theorem: if $(z - 1)X(z)$ has no poles outside the unit circle, then

$$\lim_{N \rightarrow \infty} x[N] = \lim_{z \rightarrow 1} (z - 1)X(z)$$

Exercises

- using only the fact that $u[n] \iff z/(z-1)$ and the right-shift property, find the z -transforms of the signals



- use the z -transform to find the value of $\sum_{n=0}^{\infty} n(-3/2)^{-n}$

References

Reference:

- B.P. Lathi, *Linear Systems and Signals*, Oxford University Press, chapter 5 (5.1–5.3).